SPECIALIST DENTAL SERVICES



7 WIMPOLE STREET LONDON, W1G 9SN T. 020 7580 4200 F. 020 7636 6036

POST OPERATIVE IMPLANT INSTRUCTIONS AFTER IMPLANT AND BONE GRAFT SURGERY

The following is a list of general instructions for your care following your implant surgery. Please read them carefully several times as most of your questions should be answered in the surgery.

Remember, care of the mouth after surgery has an important effect on healing.

Depending on the extent, nature and the location of the surgery, swelling, discomfort, slight oozing of blood and restricted jaw movement should be expected.

Carefully following these instructions should help you get the best results from your procedure.

How can I care for my mouth on the day after implant surgery?

- ✓ Do not rinse, swab your mouth or brush your teeth on the day after implant surgery
- Starting the following day after your surgery, rinse your mouth gently for at least 1 minute with Corsodyl mouthwash 4-5 times daily until your first post-operative appointment
- ✓ Do not pull on lips or cheeks to view the surgical site. This can open the surgical site and retard healing
- ✓ Avoid smoking, alcohol or drinking through a straw, as these may disrupt the healing, promote bleeding or pain

Can I brush my teeth following implant surgery?

- ✓ You may resume routine brushing the following day <u>BUT</u> avoid the surgical site
- ✓ Be careful when you brush the rest of the mouth, please do not disturb the surgical area.

What happens if I have bleeding?

- ✓ It is not unusual to have slight oozing for 24 hours and periodic break-through bleeding for 10 days after an oral surgery procedure. A drop or two of blood mixed with saliva may seem like more blood than there really is
- ✓ Avoid spitting or rinsing your mouth on the day of the surgery

Will the surgical site be swollen?

- You may experience swelling after surgery. The swelling peaks in appearance in 2-3 days and takes up to a week to 10 days to return to normal. Applying ice packs for the first 24 hours, with 20 minutes on and off intervals and elevating your head may help decrease swelling. If you experience swelling several weeks after your surgery, please make an appointment to contact us
- ✓ Sleep with couple of extra pillows to keep your head elevated at night for the first three nights after surgery

Am I going to have discomfort following implant surgery?

✓ A certain amount of discomfort must be expected with any surgery and medication usually will be given to you to make feel comfortable

Can I eat and drink as normal after the implant surgery?

Avoid eating until numbness is worn out. Eat whatever is comfortable. You may require soft or liquidated foods for 2-7 days. Avoid foods that require hard chewing. It is very important to maintain good dietary intake.

Can I carry on my daily activities?

✓ You should rest as much as possible for at least 24 hours after implant surgery

Do I need antibiotics?

You may be prescribed antibiotics depending on the surgery. It is very important that your antibiotics (if prescribed) be taken as directed until all are used to prevent the possibility of infection. If you develop an adverse reaction (rash or itching), discontinue the antibiotic and contact us immediately

I am wearing dentures. Can I continue wearing my dentures following implant surgery?

✓ If you wear a complete or partial denture, it is very important to follow the instructions regarding its use after surgery. It is best not to wear your denture for up to a week after surgery. Your denture will need modification before wearing is resumed

What will happen to my stitches?

✓ Stitches are placed to close the surgical site. You should not disturb the surgical area. Sutures are usually removed after 1 week after surgery

Is smoking bad for implants?

✓ Today it's known that smokers have a higher risk to loose implant than non-smokers. Irritants from smoking affect the normal healing of the gum and bone tissues of the mouth

What if I had a bone graft and/or guided bone regeneration?

✓ If you have had a bone graft, it is important to prevent any kind of pressure on the grafts. If the graft becomes exposed, or you are concerned, contact us to schedule an appointment

If you have any concerns (severe pain, swelling), please call us on 0207 580 4200 or Dr Shahdad 07813191855 or email: smile@specialistdentalservices.com