

# Application Rate Sheet

black<sup>leaf</sup>earth  
humizen

Alpha<sup>drop</sup>  
PLUS

Crop	Application	Instructions
Cereals, Oilseeds, Corn, Soybean, Fruit & Vegetable	Soil Application	Apply in band applications, in furrow, or injected into irrigation water at rates of 1–4 quarts / ac per application, up to 5 times per season, from preplant until fruit set.
	Foliar Application	Apply at rates of 1 – 2 pints / ac per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from preplant until fruiting. Caution – applications made after fruit set may stain fruit.
Tree Fruits & Vines	Soil Application	Apply broadcast or band applications, or injected into irrigation water, at rates of 1– 4 quarts per application, up to 5 times per season. Applications may be made at any point in the growing season, until & including fruit set.
	Foliar Application	Apply at rates of 1 – 2 pints / ac per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from preplant until fruiting. Caution – applications made after fruit set may stain fruit.

# Application Rate Sheet

black<sup>leaf</sup>earth  
humizen



Crop	Application	Instructions
Cereals, Oilseeds, Corn, Soybean, Fruit & Vegetable	Seed Treatment	Apply Phos-Excel Plus at a rate of 1 gal / ac, applied as a coating to appropriate amount of seed. Dilute with water as necessary prior to coating. Phos-Excel Plus is compatible with fungicidal & insecticidal seed treatments. Planting should occur within 48 hrs of dilution or blending.
	Band Applications	Apply in furrow (seed row), during seed-bed preparation, or in fertilizer bands at a rate of 1 gal / ac during or prior to planting.
	Irrigation Applications	Inject into irrigation water at rates of 1 – 4 quarts / ac. Initial application should occur within 7 days of planting. Make subsequent applications as desired at 7 – 14-day intervals, to a max of 10 applications per growing season.
Tree Fruits & Vines	Soil Application	Apply broadcast or band applications, or injected into irrigation water, at rates of 1- 4 quarts per application, up to 5 times per season. Applications may be made at any point in the growing season, until & including fruit set.
	Foliar Application	Apply at rates of 1 – 2 pints / ac per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from preplant until fruiting. Caution – applications made after fruit set may stain fruit.

# Application Rate Sheet



Crop	Application	Instructions
<b>Cereals, Oilseeds, Corn, Soybean Annual Fruit &amp; Vegetable, Forage Crops, Hemp / Cannabis</b>	Broadcast Applications	Apply Magna Plus pre-plant or pre-emergence on annual crops at rates of 20 – 75 lb / acre as a standalone product or blended with compatible dry fertilizers.
		Apply Magna Plus prior to growing season or between cuttings on perennial forage crops at rates of 20 – 75 lb / acre as a standalone product or blended with compatible dry fertilizers.
	Band Applications	Apply Magna Plus pre-plant or with fertilizers during planting at rates of 5 – 20 lb / acre as a standalone product or blended with compatible fertilizers.
		Apply Magna Plus in-furrow at rates of 5 – 20 lb / acre.
<b>Tree Fruits &amp; Vines</b>	Broadcast Applications	Apply Magna Plus prior to growing season at rates of 20 – 75 lb / acre, as a standalone application or blended with compatible dry fertilizers. Additional applications may be made as desired throughout the season. For best results, apply prior to fruit set. Apply Magna Plus prior to growing season at rates of 20 – 75 lb / acre, as a standalone application or blended with compatible dry.