## **Sprayable Body Lotion**

Reference: TR 52-11-01A

INCI name	%		Function
PHASE A			
Water (Aqua)	up to 100.00		
Glycerin	3.00		Humectant
PHASE B			
Sodium Polyacrylate	0.30		Thickener
PHASE C			
Polyglyceryl-4 Laurate (and) Polyglyceryl-6 Laurate- <b>PolyAquol™ LW</b>	3.00		Emulsifier
Oryza Sativa Bran Oil	2.50		Emollient
Gossypium Herbaceum Seed Oil	2.50		Emollient
Caprylic/Capric Triglyceride	3.00		Emollient
Octyldodecyl Oleate (and) Octyldodecyl Stearoyl Stearate (and) Polyhydroxystearic Acid (and) Octyldodecanol- innollient™ LO	2.00		Emollient
PHASE D			
Preservative	a.n.		Preservative
Parfum	a.n.		Parfum
VISCOSITY after 24h Brk, RVDV-I Prime, Sp 03, 20 rpm pH (100%)			1200 - 1800 cps 5.0 - 7.0

## Procedure:

- Prepare phase A and phase C separately and stir them at room temperature (25°C - 30°C);
- Disperse phase B in the phase A using a suitable dispersion unit;
- Add phase C to phase A+B, under agitation and homogenize for a few minutes;
- Add phase D, under gentle stirring;
- Adjust pH, if required.