

Getting started with FYCOMPA®

# SEIZE YOUR TOMORROWS

Convulsive seizure control is possible\*



\*In clinical trials, some patients taking FYCOMPA experienced fewer seizures. Individual results may vary.

## What is FYCOMPA (perampanel)?

FYCOMPA is a prescription medicine used:

- alone or with other medicines to treat partial-onset seizures with or without secondarily generalized seizures in people with epilepsy aged 4 and older
- with other medicines to treat primary generalized tonic-clonic seizures in people with epilepsy aged 12 and older

## Selected Safety Information

FYCOMPA may cause serious mental (psychiatric) problems. Tell your healthcare provider right away if you have any new or worsening mental problems while taking FYCOMPA.

Please see **Important Safety Information** on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa**<sup>™</sup>  
(perampanel) TABLETS 2•4•6•8•10•12mg  
ORAL SUSPENSION 0.5mg/mL 

# YOUR TREATMENT WITH FYCOMPA® STARTS HERE

Whether you've been recently diagnosed with epilepsy or have tried multiple treatment options before, you're looking for seizure control. You and your doctor have decided FYCOMPA might be able to help. This brochure is here to provide education and support as you begin treatment with FYCOMPA. Read on to learn about epilepsy, how FYCOMPA may help you, and what resources are available.

You're here because you want to live life on your terms. You may even have hopes for seizure freedom. With the right treatment and a healthy lifestyle, you can work toward the possibility of convulsive seizure control.

## Selected Safety Information

### What is the most important information I should know about FYCOMPA?

#### 1. FYCOMPA may cause mental (psychiatric) problems, including:

- new or worse aggressive behavior (including homicidal behavior), hostility, anger, anxiety, or irritability
- being suspicious or distrustful (believing things that are not true)
- seeing objects or hearing things that are not there
- confusion
- difficulty with memory
- other unusual or extreme changes in behavior or mood

Tell your healthcare provider right away if you have any new or worsening mental problems while taking FYCOMPA.

# TABLE OF CONTENTS

Learn About Epilepsy	6
What Is Epilepsy?.....	8
Living With Epilepsy.....	8
How FYCOMPA® Can Help	14
FYCOMPA Treats Different Types of Seizures.....	16
FYCOMPA Can Reduce Seizures.....	18
Convenient Once-Daily Dosing.....	20
How to Take FYCOMPA Liquid.....	22
Important Safety Information.....	22
Get the Support You Need	26
The FYCOMPA® Instant Savings Card Program.....	28
Connect With Others and Get Support.....	30
Eisai's Commitment to Epilepsy.....	30

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL



LEARN ABOUT  
EPILEPSY

# WHAT IS EPILEPSY?

Epilepsy is a condition of the brain that causes seizures. The brain is a complex system of nerves and cells (and a lot of other parts too). These all work together to communicate signals to different parts of the body, telling them what to do. Everything from eating, talking, and walking is controlled by and depends on these signals working right.

In people with epilepsy, **these signals become overloaded and confused**. Too many signals, or the wrong type of signals being sent at once, are what cause a seizure.

## WHAT HAPPENS DURING A SEIZURE?

How you **MOVE, ACT, or FEEL** can change during a seizure.

During a seizure you **MIGHT GET CONFUSED** and **MAY NOT BE AWARE OF WHAT'S GOING ON** around you.

## You are not alone

Millions of people in the United States and worldwide have epilepsy.

**3 MILLION**  
ADULTS LIVING WITH  
EPILEPSY IN THE US

**470,000**  
CHILDREN LIVING WITH  
EPILEPSY IN THE US

**65 MILLION**  
PEOPLE WORLDWIDE  
LIVING WITH EPILEPSY



# LIVING WITH EPILEPSY

As you may already know, living with epilepsy means it's important to consider your overall health in addition to managing your seizures. You should always take your medicine as prescribed by your doctor. It's also important to take care of your body and practice self-care. This is critical for your physical and emotional health.

Remember to take your medicine as prescribed by your doctor.  
**MISSING EVEN 1 DOSE CAN HAVE A NEGATIVE IMPACT ON YOUR HEALTH.**



## Some tips to stay healthy



Everyone needs time to wind down and relax. Taking it easy is important.



Drinking alcohol—and skipping doses of your medication to do so—can be dangerous to your health.



Think about what you're putting into your body. Are you maintaining a well-balanced diet? Consider the many benefits of eating healthy.



Stress can come on quickly and have damaging effects on your brain and body. Think of ways to unwind and focus on what's important to you.

**FINDING THE RIGHT TREATMENT OPTION FOR YOUR EPILEPSY IS IMPORTANT.**  
But it's just 1 piece of the puzzle. Staying committed to your health, as mentioned above, can also help.

## LIVING WITH EPILEPSY

---

Unfortunately, seizures can happen even if you are taking medicine. You may have heard these referred to as **breakthrough seizures**. Sometimes they occur because your medicine is no longer working for you. Often times, these seizures happen when a dose is missed because the medicine does not stay in the body for a long time.

But life is hectic and we're only human, so missed doses can happen. It's important to plan ahead as best you can and take your medicine as prescribed to try to avoid experiencing a seizure while on treatment.





HOW  
**FYCOMPA**<sup>®</sup>  
CAN HELP

Please see Important Safety Information on pages 22 to 23  
and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa**<sup>™</sup>  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL 

# FYCOMPA® TREATS DIFFERENT TYPES OF SEIZURES



## PARTIAL

### PARTIAL-ONSET SEIZURES (ALSO KNOWN AS "FOCAL")

Happen in one part of the brain


- Usually last less than 2 minutes
- Experiences can vary depending on where the seizure occurs in the brain
- Stiffening or jerking movements
- Changes in smell, taste, and hearing
- Strange sensations in the body can occur or impact how people think or experience events
- Sometimes memory is lost

### Selected Safety Information

FYCOMPA may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call your healthcare provider right away if you have any new or worsening symptoms.

FYCOMPA® is a prescription medicine used:

- to treat **partial-onset seizures** with or without **secondarily generalized seizures** in people with epilepsy who are 4 years of age and older
- with other medicines to treat **primary generalized tonic-clonic seizures** in people with epilepsy who are 12 years of age and older



## CONVULSIVE

### SECONDARILY GENERALIZED SEIZURES

Start on one side of the brain and spread to the other

### PRIMARY GENERALIZED TONIC-CLONIC SEIZURES (ALSO KNOWN AS "GRAND MAL")

Can start on both sides of the brain at the same time

- Usually last 1 to 3 minutes
- Happen without warning
- Movements can be strong or forceful
- Loss of consciousness, falling to the floor
- Breathing can be temporarily impaired
- No memory of the event happening

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

# IT'S TIME TO SEIZE YOUR TOMORROWS

## FYCOMPA® may help control your seizures

The studies used to gain FDA approval of FYCOMPA studied how effective FYCOMPA was at reducing certain types of seizures.

Some patients 12 years of age and older treated with FYCOMPA\* experienced:



A decrease in the number of seizures they had and how often seizures happened

Not all patients had the same results with FYCOMPA

\*FYCOMPA plus 1 to 3 antiepileptic drugs.

Based on this research:

Similar results are expected in patients with partial-onset seizures when FYCOMPA is used alone or in children 4 years of age or older

### Selected Safety Information

#### What should I avoid while taking FYCOMPA?

Do not drive, operate heavy machinery, or do other dangerous activities until you know how FYCOMPA affects you. FYCOMPA may make you dizzy, sleepy, or tired. Do not drink alcohol or take other medicines that make you sleepy or dizzy until you talk to your healthcare provider. FYCOMPA taken with alcohol or medicines that cause sleepiness or dizziness may make your sleepiness or dizziness worse. FYCOMPA when taken with alcohol may also make your mood worse, increase anger, confusion, and depression.

## Is seizure freedom possible for me?



Clinical studies provide information about the safety and effectiveness of a medicine.

The main purpose of the FYCOMPA® studies was measuring the number of seizures and how often the seizures happened. Sometimes researchers look for results that were not part of the main purpose of the study. In the FYCOMPA studies, researchers also looked for patients who achieved seizure freedom (meaning patients had no seizures) over the course of 13 weeks. These results were measured either during the study or after the study ended.



While measuring seizure freedom was not the main purpose of the studies, results from these separate reviews of the data showed that some patients achieved seizure freedom while taking FYCOMPA.



Because measuring seizure freedom was not the main purpose of the FYCOMPA studies, the data are not considered to be significant. This means the data are not meaningful enough to be able to draw conclusions about these results.

**While some patients experienced seizure freedom with FYCOMPA, everyone reacts differently to medicine and may experience different results.**

**TALK TO YOUR DOCTOR TO LEARN MORE ABOUT FYCOMPA**

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

# CONVENIENT ONCE-DAILY DOSING

Living with epilepsy already has its challenges. The simple administration of FYCOMPA® fits into your daily routine.



FYCOMPA comes in small tablet or liquid form.



**FYCOMPA HAS A LONG HALF-LIFE**

Half-life describes how long a medicine stays in your body. Half-life is important in epilepsy treatment because medicines with longer half-lives stay in your body at steadier levels for a longer time even in the event you miss a dose. Talk to your doctor if you miss one or more of your doses or if you have questions about what a long half-life means to you. Remember to always take your medication as prescribed by your doctor.

## Selected Safety Information

**Do not stop FYCOMPA without first talking with your healthcare provider.** Stopping suddenly can cause serious problems and can cause you to have seizures more often.

## FYCOMPA® is taken once daily, whenever you go to bed



**START FYCOMPA LIQUID OR TABLET**

Depending on what type of seizures you are experiencing, FYCOMPA can be taken alone or with other medicines. Your doctor will first start you on a low dose. Then, your doctor will slowly increase your dose to be sure you get the amount of FYCOMPA that's right for you. Your dose could be increased as slowly as every 2 to 4 weeks, but no more than once a week, depending on how FYCOMPA works for you.



**TAKE FYCOMPA AT BEDTIME**

You should take FYCOMPA every night at bedtime, whenever that may be. Sometimes life can get hectic so your bedtime may be different from night to night and that's OK. **It won't change the way FYCOMPA works.**



**MONITOR FOR ANY REACTIONS**

If you notice any reactions to FYCOMPA, it's important to let your doctor know right away. Your doctor may lower or discontinue your dose depending on how you react to FYCOMPA.

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL



# HOW TO TAKE FYCOMPA<sup>®</sup> LIQUID

- YOU WILL NEED:**
- Liquid formulation bottle
  - Bottle adapter
  - Dosing syringe

## STEP 1

Take the Fycompa bottle, bottle adapter, and syringe out of the box.


## STEP 2

Shake the bottle well—like you would shake a container of salad dressing—before each use.



## STEP 3

Take the cap off the bottle. Put the bottle adapter into the bottle by pressing it down. After the bottle adapter is in place, you can't take it off. **See A**



## STEP 4

Check the dose in milliliters (mL) of Fycompa liquid that your doctor prescribed. Find this number on the syringe. **See B**

## STEP 5

Push the plunger of the syringe all the way down. While the bottle is upright, put the syringe into the bottle through the opening in the bottle adapter. **See C**



## STEP 6

Hold the syringe in place, and turn the bottle and syringe upside down. Pull the plunger on the syringe to reach the dose your doctor prescribed (the amount of Fycompa medicine in Step 4). If you see air bubbles in the syringe, fully push in the plunger so the medicine flows back into the bottle. Then, pull the plunger on the syringe to reach the dose your doctor prescribed. Measure the mLs of Fycompa liquid from the white line at the end of the plunger, not the black line.



If your prescribed dose is more than 20 mL, go to Step 7. If your dose is 20 mL or less, skip Step 7 and go to Step 8. **See D**

## STEP 7


If your dose is more than 20 mL, you can get the right amount of Fycompa by using the same syringe. Draw the first 20 mL into the syringe and slowly squirt into the corner of your mouth (Steps 8 to 9). Then, draw the remaining amount of medicine in the same syringe (Steps 4 to 6) and take Fycompa liquid (Steps 8 to 9).

## STEP 8

Turn the bottle right-side up and take the syringe out of the bottle adapter.

## STEP 9


Push down on the plunger to slowly squirt Fycompa<sup>®</sup> directly into the corner of your mouth. If your dose is more than 20 mL, slowly squirt Fycompa from the syringe into the corner of your mouth. Then repeat Steps 7 to 8 and slowly squirt Fycompa from the same syringe into the corner of your mouth. **See E**



## STEP 10

Rinse the syringe with tap water after each use. To do this:

- Fill a cup with water
- Put the tip of the syringe in the water. Pull the plunger back to draw water from the cup into the syringe
- Push down the plunger to release water into the sink. **See F**



## STEP 11

Put the cap on the bottle and twist tightly. Leave the bottle adapter on. The cap will fit over the bottle adapter.

Please see **Important Safety Information** on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

# HOW TO STORE FYCOMPA liquid

- ✓ Put the cap on the bottle. Twist it on tightly
- ✓ Store Fycompa liquid below 86°F (30°C). Do not freeze
- ✓ Replace the cap tightly after opening
- ✓ Use Fycompa liquid within the first 90 days of opening the bottle
- ✓ Do not use Fycompa liquid that has been in an open bottle for more than 90 days. Safely throw out any Fycompa liquid that has not been used within 90 days of being opened

**Fycompa<sup>™</sup>**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

## Important Safety Information

### What is the most important information I should know about FYCOMPA®?

1. **FYCOMPA may cause mental (psychiatric) problems, including:**
  - new or worse aggressive behavior (including homicidal behavior), hostility, anger, anxiety, or irritability
  - being suspicious or distrustful (believing things that are not true)
  - seeing objects or hearing things that are not there
  - confusion
  - difficulty with memory
  - other unusual or extreme changes in behavior or mood

Tell your healthcare provider right away if you have any new or worsening mental problems while taking FYCOMPA.

2. **Like other antiepileptic drugs, FYCOMPA may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call your healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:**
  - thoughts about suicide or dying
  - new or worse depression
  - feeling agitated or restless
  - trouble sleeping (insomnia)

- acting aggressive, being angry, or violent
- an extreme increase in activity and talking (mania)
- attempt to commit suicide
- new or worse anxiety
- panic attacks
- new or worse irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

### How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes especially sudden changes in mood, behaviors, thoughts, or feelings
- Keep all follow-up visits with your healthcare provider as scheduled

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

**Do not stop FYCOMPA without first talking with your healthcare provider.** Stopping suddenly can cause serious problems and can cause you to have seizures more often.

### Before taking FYCOMPA, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had depression, mood problems, aggressive or hostile behavior (for example, homicidal behavior), suicidal thoughts or behavior, or other psychiatric problems
- have liver or kidney problems
- drink alcohol
- have abused prescription medicines, street drugs, or alcohol in the past
- are pregnant or plan to become pregnant. It is not known if FYCOMPA will harm your unborn baby. If you become pregnant while taking FYCOMPA, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry (1-888-233-2334)
- are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you take FYCOMPA and to decide if you will take FYCOMPA or breastfeed. You should not do both.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking FYCOMPA® with certain other medicines can cause side effects or reduce either drug's benefit. These medicines include: birth control, carbamazepine, phenytoin, oxcarbazepine, rifampin, and St. John's Wort.

### What should I avoid while taking FYCOMPA?

Do not drive, operate heavy machinery, or do other dangerous activities until you know how FYCOMPA affects you. FYCOMPA may make you dizzy, sleepy, or tired. Do not drink alcohol or take other medicines that make you sleepy or dizzy until you talk to your healthcare provider. FYCOMPA taken with alcohol or medicines that cause sleepiness or dizziness may make your sleepiness or dizziness worse. FYCOMPA when taken with alcohol may also make your mood worse, increase anger, confusion, and depression.

**What are the possible side effects of FYCOMPA?** FYCOMPA may cause other serious side effects, including:

- **Dizziness, vertigo (sense of spinning), and problems walking normally.** You may have problems walking normally if you are unsteady because you feel dizzy. These symptoms can increase when your dose of FYCOMPA is increased. Your risk of feeling dizzy and having problems walking normally may be higher if you are elderly.
- **Sleepiness and tiredness**
- **Increased risk of falls.** Taking FYCOMPA can increase your chance of falling. These falls can cause serious injuries. Your risk of falling may be higher if you are elderly.
- **A serious allergic reaction that may affect your skin or other parts of your body such as your liver, kidneys, heart, or blood cells.** This allergic reaction can be life-threatening and can cause death. Call your healthcare provider right away if you have:
  - a skin rash, hives
  - fever or swollen glands that do not go away
  - swelling of your face
  - shortness of breath, swelling of the legs, yellowing of the skin or whites of the eyes, or dark urine

The most common side effects of FYCOMPA include:

- dizziness
- sleepiness
- tiredness
- irritability
- falls
- nausea and vomiting
- weight gain
- vertigo (sense of spinning)
- problems walking normally
- problems with muscle coordination
- headache
- bruising
- abdominal pain
- anxiety

FYCOMPA is a controlled substance (CIII) because it can be abused or lead to drug dependence. Keep FYCOMPA in a safe place to protect it from theft and never give it to anyone else because it may harm them. Selling or giving away FYCOMPA is against the law.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

Please see [Prescribing Information](#) and [Medication Guide](#).



GET THE  
**SUPPORT**  
YOU NEED

Please see Important Safety Information on pages 22 to 23  
and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5mg/mL

# THE FYCOMPA® INSTANT SAVINGS CARD PROGRAM

PAY AS LITTLE AS \$10 PER MONTH\* FOR FYCOMPA®

Once you've been prescribed FYCOMPA, you can receive your savings card through your doctor or by visiting [FYCOMPA.com](https://www.fycompa.com) and clicking the "Register or Activate now" link. You can also activate the card by calling 1-855-347-2448.

- Bring your card to the pharmacy
- Show your savings card to the pharmacist each time you fill your prescription

The back of the card has details for the savings program. This will help ensure that the offer applies to you. Eligible commercially insured patients may pay as little as \$10 per month with a maximum savings of \$1,300 per year.

Eligible cash patients can receive up to \$60 per prescription for a maximum savings of \$720 per year.

Ask your doctor or pharmacist if you are not sure whether the offer applies to you.



## FYCOMPA® Patient Assistance Program

In addition to the FYCOMPA® Instant Savings Card, this program provides information that may help you pay for your medicine.

If you qualify, you may be able to get FYCOMPA® at no cost and receive other resources.

Call **1-855-347-2448** or visit [EisaiReimbursement.com](https://EisaiReimbursement.com) to learn more.

Eisai is **COMMITTED TO SUPPORTING PATIENTS** by providing information that may help you get access to the treatment you need

\*Restrictions apply. Not available to patients enrolled in federal or state healthcare programs, including Medicare, Medicaid, Medigap, VA, DoD, or TRICARE. See <https://www.fycompa.com/savings-card> for complete terms and conditions.

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

## CONNECT WITH OTHERS AND GET SUPPORT

Living with epilepsy can be stressful, but you don't have to face it alone. Keep an open channel of communication with your doctor in case you have any questions or concerns about treatment with FYCOMPA®.

MORE THAN **200,000** PEOPLE WORLDWIDE  
HAVE BEEN PRESCRIBED FYCOMPA<sup>\*†</sup>

It can also help to talk to other people who have epilepsy. You can learn a lot from them and they can learn a lot from you, too. You can start by hearing from patients like you who wanted to share their story at [FYCOMPA.com](https://www.fycompa.com).

\*Worldwide figure from 2012 through July 2018. 30,366 patients prescribed FYCOMPA in the United States.

†Across different indications.

### Other helpful resources:

**Advancing Epilepsy Care**  
advancingepilepsycare.com

**Centers for Disease Control and Prevention (CDC)**  
CDC.gov/epilepsy

**Citizens United for Research in Epilepsy (CURE)**  
CUREepilepsy.org

**National Association of Epilepsy Centers (NAEC)**  
NAEC-epilepsy.org

**The Epilepsy Foundation**  
Epilepsy.com

**North American Antiepileptic Drug Pregnancy Registry**  
AEDpregnancyregistry.org

IF YOU ARE CARING  
FOR SOMEONE  
WITH EPILEPSY

Visit [FYCOMPA.com](https://www.fycompa.com) for information specifically for caregivers.



Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).

**Fycompa**<sup>™</sup>  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

# EISAI'S COMMITMENT TO EPILEPSY

Eisai is a pharmaceutical company committed to putting patients and families first, and is driven by a deeply rooted history of helping patients and caregivers. Eisai has studied multiple medicines for the treatment of epilepsy.

## Community outreach

Eisai is actively involved in the community and supporting foundations that help spread awareness about epilepsy. Find out more about Eisai's commitment to patients with epilepsy by visiting [us.eisai.com](https://us.eisai.com).



**Fycompa™**  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL

FYCOMPA®:

# SEIZE YOUR TOMORROWS

Interested in learning more? [Visit FYCOMPA.com](http://Visit FYCOMPA.com)



## Selected Safety Information

The most common side effects of FYCOMPA include: dizziness; sleepiness; tiredness; irritability; falls; nausea and vomiting; weight gain; vertigo (sense of spinning); problems walking normally; problems with muscle coordination; headache; bruising; abdominal pain; anxiety.

Please see Important Safety Information on pages 22 to 23 and [Prescribing Information](#) and [Medication Guide](#).



FYCOMPA® is a registered trademark of Eisai R&D Management Co., Ltd., licensed to Eisai Inc.  
©2019 Eisai Inc. FYCO-US2958 June 2019

**Fycompa**<sup>™</sup>  
(perampanel) TABLETS 2•4•6•8•10•12 mg  
ORAL SUSPENSION 0.5 mg/mL